

ogy, Southwest Texas State University, San Marcos, Texas 78666, USA (e-mail: mg45447@swt.edu), **JAMES R. DIXON**, Texas Cooperative Wildlife Collection, Texas A&M University, College Station, Texas 77843, USA, and **MICHAEL R. J. FORSTNER**, Department of Biology, Southwest Texas State University, San Marcos, Texas 78666, USA (e-mail: mf11@swt.edu).

ODONTOPHRYNUS LAVILLAI. ARGENTINA: SAN JUAN: Valle Fértil Department, 137.5 km from San Juan (city) on Las Tumanas, Las Tumanas River (30°52'S, 67°19'W, 750 m elev). 25 February 2001. J. H. Villavicencio and M. G. Cánovas. Herpetological Collection Instituto y Museo de Ciencias Naturales, Universidad Nacional de San Juan, Argentina (IMCN-UNSJ 545). Verified by Juan Carlos Acosta. First province record; extends the known range ca. 350 km SW from the nearest locality of Sierra de Guasayán, Choya Department, Santiago del Estero Province, the type locality (Ceí 1985, Cuad. Herpetol. 1[5]:1–13).

Submitted by **JOSE HECTOR VILLAVICENCIO** and **MARIA GABRIELA CANOVAS**, Departamento de Geofísica y Astronomía, F.C.E.F. y N. Universidad Nacional de San Juan, Avenida España 400 (N), Caixa Postal 5400, San Juan, Argentina; e-mail (MGC): dgocanovas@sinectis.com.ar.

PSEUDACRIS CRUCIFER CRUCIFER (Northern Spring Peeper). USA: WISCONSIN: DODGE CO: SE 1/4 Sec. 5, T13N, R13E, calling from south of Lake Emily. 20 April 2001. Jack and Holly Bartholmai. MPM 32932. Verified by Robert W. Henderson. New county record (Casper 1996, Geographic Distributions of the Amphibians and Reptiles of Wisconsin. Publ. Milwaukee Pub. Mus., Milwaukee. 87 pp.; Vogt 1981, Natural History of Amphibians and Reptiles of Wisconsin. Publ. Milwaukee Pub. Mus., Milwaukee. 205 pp.). Verifies reported observations from Dodge County Wisconsin Frog and Toad Survey Routes (Mossman et al. 1998, *In* M. J. Lannoo [ed.], Status and Conservation of Midwestern Amphibians, pp. 169–198, Univ. Iowa Press, Iowa City).

Submitted by **GARY S. CASPER**, Section of Vertebrate Zoology, Milwaukee Public Museum, 800 West Wells Street, Milwaukee, Wisconsin 53233, USA; e-mail: gsc@mpm.edu.

PSEUDACRIS STRECKERI (Strecker's Chorus Frog). USA: ARKANSAS: PULASKI CO: (N34.81341°, W92.40218°), Two Rivers Park. 12 March 2001. K. J. Irwin and L. K. Irwin. Arkansas State University Museum of Zoology, Herpetological Collection (ASUMZ 26099). Verified by Stanley Trauth. Adult male calling in flooded park field. Syntopic species associates were *Pseudacris crucifer* and *Pseudacris feriarum*. First record for county and represents the most southeasterly record for the species in the state. (Trauth, Robison, and Plummer, ms. in prep.).

Submitted by **KELLY J. IRWIN**, Arkansas Game & Fish Commission, 915 East Sevier Street, Benton, Arkansas 72015, USA (e-mail: kirwin@agfc.state.ar.us), and **LISA K. IRWIN**, 103 Brooke Drive, Perryville, Arkansas 72126, USA.

RANA GLANDULOSA (Rough-sided Frog). INDONESIA: NATUNA ISLANDS: Bunguran Island (4°00'N, 108°15'E). August 1928. F. N. Chasen. Raffles Museum of Biodiversity Research Zoological Reference Collection (ZRC.1.613, adult female, SVL 84.7 mm). Verified by Kelvin K. P. Lim. New island record; not

previously recorded from the Natunas (Van Kampen 1923, The Amphibia of the Indo-Australian Archipelago. E. J. Brill, Leiden, xii + 304 pp.).

Submitted by **TZI MING LEONG**, Department of Biological Sciences, National University of Singapore, Singapore 119260; e-mail: scip0132@nus.edu.sg.

RANA PALUSTRIS (Pickerel Frog). USA: ARKANSAS: PHILLIPS CO: (N34.57202°, W90.59502°), SE 1/4, Sec. 29, R5E, T1S, St. Francis National Forest. 3 March 2001. Ron Bonett, Carol Spence, L. K. Irwin, and K. J. Irwin. Arkansas State University Museum of Zoology, Herpetological Collection (ASUMZ 26098). Verified by Stanley Trauth. Subadult found under leaves on gravel substrate spring outflow at base of eastern versant of Crowley's Ridge. Syntopic species associates were *Acris crepitans*, *Rana clamitans*, and *Rana sphenoccephala*. First record for Phillips County (Trauth, Robison, and Plummer, ms. in prep.).

Submitted by **RON BONETT**, Department of Biology, University of Texas at Arlington, Arlington, Texas 76019, USA (e-mail: desmog_2000@yahoo.com), **KELLY J. IRWIN**, Arkansas Game and Fish Commission, 915 East Sevier Street, Benton, Arkansas 72015, USA (e-mail: kirwin@agfc.state.ar.us), **CAROL SPENCE**, Department of Biology, University of Texas at Arlington, Arlington, Texas 76019, USA, and **LISA K. IRWIN**, 103 Brooke Drive, Perryville, Arkansas 72126, USA.

RANA SYLVATICA (Wood Frog). USA: TENNESSEE: GRUNDY CO: Savage Gulf State Natural Area (35°28'01"N, 85°36'02"W). 3 May 2001. Stephen T. Samoray and Kurt J. Regester. Austin Peay State University Museum (APSU Color Photograph 3273). Verified by A. Floyd Scott. One adult found on Connector Trail ca. 1 km south of Hobbs cabin. New county record (Redmond and Scott 1996, Austin Peay State Univ. Center Field Biol. Misc. Publ. 12:1–94); extends range ca. 45 km SW of previously recorded limit in Bledsoe County.

Submitted by **STEPHEN T. SAMORAY** and **KURT J. REGESTER**, Department of Biology, Middle Tennessee State University, Murfreesboro, Tennessee 37132, USA.

SCINAX NASICUS. ARGENTINA: CÓRDOBA: Río Cuarto (33°08'S, 64°22'W), 434 m elev. 9 March 2001. N. E. Salas. Herpetological Collection of Universidad Nacional de Río Cuarto, Ecología. (UNRC-ECO 690). Verified by R. Martori. Previously recorded in Argentina from Jujuy, Salta, Formosa, Chaco, Misiones, Córdoba, Santiago del Estero, Tucumán, Corrientes Santa Fe, Entre Ríos, and Buenos Aires (Ceí 1980, Amphibians of Argentina. Monit. Zool. Ital. [N. S.], Monogr. 2:1–609; Lavilla et al. 2000. *In* Lavilla et al. [eds.], Categorización de los Anfibios y Reptiles de la República Argentina, pp. 11–34. Asociación Herpetológica Argentina). In Córdoba, was reported from the northeastern departments of Sobremonte, Río Seco, Tulumba, and San Justo (di Tada et al. 1996. *In* I. E. di Tada and E. H. Bucher [eds.], pp. 191–213. Ed. UNRC), and from the eastern departments of General San Martín and Unión (Rozzi Giménez and Gallego 1998, Cuad. Herpetol. 14[2]:169). First report for southern Córdoba and extends distribution 130 km south.

Submitted by **NANCY E. SALAS** and **ISMAEL E. DI TADA**, Departamento de Ciencias Naturales, Facultad de Ciencias Exactas,